

The Panda Math Tutor

www.thepandamathtutor.com

Algebraic Fractions

IGCSE Mathematics

Prepared by

Nikolas Agisilaou

Quality Mathematics Education for Students

Contents

1	Introduction to Algebraic Fractions	2
2	Simplifying Algebraic Fractions	2
3	Multiplying and Dividing Algebraic Fractions	5
4	Adding and Subtracting Algebraic Fractions	8
4.1	Same Denominators	8
4.2	Different Numerical Denominators	9
4.3	Different Algebraic Denominators	10
5	Solving Equations with Algebraic Fractions	12
6	Forming and Solving Problems Using Algebraic Fractions	15
6.1	Worked Problems	15
7	Further Techniques	16
7.1	Fractions with Repeated Factors	17
7.2	Simplifying Complex Fractions	17
8	Summary	18
9	Additional Resources	19
9.1	Tips for Success in IGCSE Exams	19
9.2	Common Exam Question Types	20

1 Introduction to Algebraic Fractions

Note

Algebraic fractions are fractions in which the numerator, denominator, or both contain algebraic expressions. They obey exactly the same rules as numerical fractions — the key additional skill is *factorising polynomials* so that common factors can be identified and cancelled.

In this chapter, we will explore:

- What algebraic fractions are and how to simplify them
- Multiplying and dividing algebraic fractions
- Adding and subtracting algebraic fractions with different denominators
- Forming and solving equations involving algebraic fractions
- Common mistakes and how to avoid them

2 Simplifying Algebraic Fractions

Definition

An **algebraic fraction** is a fraction whose numerator, denominator, or both are algebraic expressions. The following are all examples:

$$\frac{3p}{7q}, \quad \frac{x^2 - 36}{x + 6}, \quad \frac{2x^2 + 9x + 4}{x^2 + 2x - 8}$$

To **simplify** an algebraic fraction means to write it in its lowest terms by cancelling any factors that appear in both the numerator and denominator.

Important: You can only cancel *factors* (connected by multiplication), never *terms* (connected by addition or subtraction).

Warning

The most common mistake — cancelling terms instead of factors!

When the numerator and denominator are sums or differences, you *cannot* cancel individual terms across the fraction bar:

$$\frac{x + 6}{x + 9} \neq \frac{6}{9} \quad \text{WRONG} \quad (x \text{ is a term, not a factor here!})$$

Similarly, incorrect partial cancellation of a single term in a sum:

$$\frac{x^2 + 4x}{x} = x^2 + 4 \quad \text{WRONG}$$

Correct approach: factorise the numerator first, then cancel the whole factor:

$$\frac{x^2 + 4x}{x} = \frac{x(x + 4)}{x} = x + 4$$

Always factorise *completely* before cancelling anything.

Method

Simplifying Algebraic Fractions

Step 1: Factorise the numerator completely

Step 2: Factorise the denominator completely

Step 3: Cancel any factor that appears in *both* numerator and denominator

Step 4: Write the simplified result; if no common factor exists, the fraction is already in its simplest form

Tip: If you cannot find a common factor after full factorisation, the fraction cannot be simplified further.

Example

Simplify $\frac{15p^3q^2}{10p^2q^5}$.

Solution:

Handle the numerical coefficient and each variable separately:

$$\frac{15p^3q^2}{10p^2q^5} = \frac{15}{10} \cdot \frac{p^3}{p^2} \cdot \frac{q^2}{q^5} = \frac{3}{2} \cdot p \cdot \frac{1}{q^3} = \frac{3p}{2q^3}$$

Answer: $\frac{3p}{2q^3}$

Preview ends here

18 more pages are in the full pack.

Algebraic Fractions — Main Notes — €9.99

Buy the pack and the complete PDF lands in your inbox.

Personal study use only — please don't resell or share.