

The Panda Math Tutor

www.thepandamathtutor.com

Curved Graphs

Solutions & Answer Key

Full worked solutions, drawn graphs, marking scheme & answer key
IGCSE Mathematics

Prepared by

Nikolas Agisilaou

Companion to the *Student Worksheet*

Contents

4	Part IV — Complete Solutions	2
4.1	Solutions — Quadratic Graphs and Their Features	2
4.2	Solutions — The Completed Square Form	5
4.3	Solutions — Reciprocal Graphs	7
4.4	Solutions — Graphical Solutions of Quadratic Equations	10
4.5	Solutions — Cubic and Exponential Graphs	12
4.6	Solutions — Gradient of a Curve	16
4.7	Solutions — Differentiation: The Power Rule	17
4.8	Solutions — Turning Points, Tangents and Normals	19
4.9	Solutions — Mixed Review	22
4.10	Marking Scheme — Examination-Style Practice	24
5	Part V — Common Mistakes & Learning Tips	29
6	Part VI — Final Answer Key	31

Note

About this booklet. These are the complete solutions to every question in the *Curved Graphs — Student Worksheet*. Question numbers match exactly (QG1–QG7, CS1–CS7, RG1–RG7, GS1–GS6, CE1–CE6, GT1–GT4, DF1–DF9, TP1–TP7, the Mixed Review MR1–MR10, and the examination questions Q1–Q8).

Every graph the student is asked to draw is reproduced here in full, so the shape and the key points can be compared directly against their own drawing. Part V collects the common mistakes to watch for, and Part VI is a quick answer key for fast marking.

4 Part IV — Complete Solutions

Note

Reading a value *from a graph* is only ever an estimate, so answers to graphical questions are given to 1 decimal place and a small tolerance should be allowed when marking. Answers obtained by *differentiating* are exact.

4.1 Solutions — Quadratic Graphs and Their Features

Solution QG1

The direction depends only on the sign of a ; the y -intercept is the constant term c .

- (a) $a = 2 > 0$: opens **upward** (minimum). y -intercept $(0, 3)$.
- (b) $a = -1 < 0$: opens **downward** (maximum). y -intercept $(0, -6)$.
- (c) Rewriting as $y = -3x^2 + 5$, we have $a = -3 < 0$: opens **downward**. y -intercept $(0, 5)$.

Solution QG2

x	-3	-2	-1	0	1	2	3
y	5	0	-3	-4	-3	0	5

Roots: $x = -2$ and $x = 2$. **y -intercept:** $(0, -4)$.

The table is symmetrical about $x = 0$, which is the axis of symmetry.

Solution QG3

Here $a = 1$, $b = -6$, $c = 5$.

$$x = -\frac{b}{2a} = -\frac{-6}{2(1)} = 3.$$

Substituting: $y = 3^2 - 6(3) + 5 = 9 - 18 + 5 = -4$.

Vertex: $(3, -4)$. **Axis of symmetry:** $x = 3$.


Check: $y = (x - 1)(x - 5)$, so the roots are 1 and 5, whose midpoint is indeed 3. ✓

Solution QG4

(a) Building the table row by row:

x	-2	-1	0	1	2	3	4
x^2	4	1	0	1	4	9	16
$-2x$	4	2	0	-2	-4	-6	-8
-3	-3	-3	-3	-3	-3	-3	-3
y	5	0	-3	-4	-3	0	5

(b) The graph:



(c) **Vertex** $(1, -4)$; **roots** $x = -1$ and $x = 3$.

Since $x^2 - 2x - 3 = (x + 1)(x - 3)$, the roots read from the graph are exact.

Preview ends here

29 more pages are in the full pack.

Curved Graphs — Solutions — €9.99

Buy the pack and the complete PDF lands in your inbox.

Personal study use only — please don't resell or share.